EARLY PERMIAN PALAEOFLORAS FROM SOUTHERN BRAZILIAN GONDWANA: A PALAEOCLIMATIC APPROACH

MARGOT GUERRA-SOMMER AND MIRIAM CAZZULO-KLEPZIG

ABSTRACT In evaluating the parameters supplied by the taphofloras from different sedimentary facies in the Early Permian sedimentary sequences of the southern part of the Paraná basin, Brazil, it has become evident that the palaeofloristic evolution was related to palaeoecological and palaeoclimatic evolution. The homogeneous composition of Early Permian floral assemblages, which are characterized mainly by herbaceous to shrub-like plants considered to be relicts from the rigorous climate of an ice age (e.g. *Botrychiopsis plantiana*) suggest the persistence of the cold climate. The dominance of *Rubidgea* and *Gangamopteris* leaves with palmate venation associated with glossopterids with pinnate venation seems to indicate a gradual warming of climate. In roof-shales of coalbearing strata pinnate glossopterids related to *Glossopteris* are common, while *Gangamopteris* and *Rubidgea* (palmate forms) are poorly represented. The sudden enrichment of herbaceous articulata and filicoids fronds is characteristic of this stage and trunks of arborescent lycophytes become important elements. These antrocophilic paleofloras are characterized by typical elements of the "*Glossopteris* flora" associated to tree lycophytes and ferns communities. Therefore, the cool seasonal climate of Early Permian changed into the moist seasonal interval during the Artinskian-Kungurian. This climatic change was significant to the meso-hygrophytic to hygrophytic vegetation registered in roof-shale ferns of the Gondwana Southern Brazilian coalbearing strata.

Keywords: roof-shale floras, Gondwana, Southern Brazil, Paraná basin, *Glossopteris* Flora

INTRODUCTION The intracratonic Paraná basin, with a total area of 1.700.000 Km² (1.100.000 Km² of which correspond to Brazil) is situated in east central South America. It contains Paleozoic, Mesozoic and locally, also Cenozoic sedimentary rocks. In southern Brazil the major coal-bearing strata occur in the intracratonic Paraná basin, in isolated coalfields located along the margin of the Precambrian shield, from the southernmost part of the state of Rio Grande do Sul, through Santa Catarina to the north of the state of Paraná. The coal measures are considered to range in age from Artinskian to Kungurian (Burjack 1978) and occur in the Rio Bonito Formation, included in the Carboniferous-Early Triassic Sequence (regressive package - Fig. 1).

Figure 1 - Location of the Gondwana Sequence in Rio Grande do Sul, southern Brazil (modified after Scherer *et al.* 1999)

Economic interest in coal seams in the Southern Brazilian Gondwana sequence led to the documentation of the megafloristic composition of the coal-bearing strata at the end of the XIX Century and at the beginning of the XX (Plant 1869, Carruthers 1869, Feistmantel 1876-1886, Zeiller 1895, Seward 1903, Arber 1905). White (1908) introduced the term "pure *Glossopteris* flora" to characterize these associations in coalfields of the states of Santa Catarina and Rio Grande do Sul. Relevant studies on megafloras from coal-bearing strata of Southern Brazil were developed since the first half of the century (Lundqvist 1919, Mendes 1952, Rigby 1970, Guerra-Sommer *et al.* 1991). These studies aim characterize the evolution of Early Permian palaeofloras of the southernmost Paraná basin, emphasizing the "antrocophilic" or roof-shale floras. These floras correspond to fossil plant association related to coal seams, but without any further implication regarding the relationship between the vegetation and coal formation (Meyen 1987).

In the present study is adapted the chronological representation

proposed by Milani *et al.* (in press) for the Paraná basin. These authors, using sequence stratigraphy, characterize six megasequences for the Paleozoic and Mesozoic intervals of the Basin limited by interregional unconformities and partially eroded. Each megasequence consist of lithological packages formally individualized as lithostratigraphic units by different authors.

PALEOFLORISTIC SUCCESSION The lower part of the Early Permian Gondwana succession in the Southern Brazil is characterized by sediments comprising mainly sandstones and diamictites, deposited in a marked periglacial climate. The megafloral remains recovered from glaciocontinental sediments related to lowlands (Milani *et al.* in press) and included into Itarare Group are assigned to Glossopteridales (*Gangamopteris obovata*, *Gangamopteris buriadica*, *Gangamopteris angustifolia*, *Rubidgea lanceolatus*, *Rubidgea obovata*, *Glossopteris indica* Schimper 1874, *Glossopteris communis*, Feistmantel 1896). *Botrychiopsis plantiana* (Carruthers) Arch. Arrondo 1971, is the second taxon in importance; on the other hand, Cordaitales (*Cordaites hislop*, Unger 1850), Coniferales (*Buriadia* sp.), Ginkgoales (*Ginkgoites eximia* Seward 1919) are complementary elements. Herbaceous Pteridophyta are rare, represented by a few preserved specimens of *Phyllothea* sp. (Acampamento Velho, Cambaf, Faxinal, Fazenda Goulart outcrops). The homogenous composition of Early Permian floral assemblages, which are dominated by foliar organs of plants with arborescent habit and shrub-like plants (*Botrychiopsis plantiana*) indicates the persistence of a cold climate (Cazzulo-Klepzig & Guerra-Sommer 1983).

With the retreat of ice caps, the glaciogene sedimentation was replaced by fluviodeltaic and estuarine facies. The taphofloras of the basal siltstones of this sequence recovered from the lower section of Papaleo outcrop (Pasqualini *et al.* 1986) are characterized by the presence of *B. plantiana* associated with articulata (*Phyllothea indica* Bunburi 1861), conifers (*Buriadia* sp.), and glossopterids with dominant pinnate venation. *Gangamopteris*, *Cordaites* and protoglossopterids (*Rubidgea*) occur as minor elements.

The increasing of herbaceous articulata in this level in relation to those of the basal Permian sequences, and evidences of foliar phytofagy point out to a climate amelioration. The depositional sequence grades upward to more restricted environment conditions that allowed the formation of peat swamps, formed in lagoons protected by barriers under retrograding regime (Holz 1998, Milani *et al.* 1997).

Detailed studies of "roof-shale" floras have revealed some vegetation heterogeneity in roof-shale floras within any one coalfield in Southern Brazilian Gondwana. Some floristic assemblages contains dominant elements of gymnospermic affinity whereas a rich pteridophytic association is registered in others.

The presence of a rich coalfield compressed flora from a "tonstein" layer associated with a coal seam in the Faxinal Mine allowed the identification of an autochthonous taphoflora characterized by the predominance of arborescent gymnospermic forms (Guerra-Sommer 1989, 1992). Fragments of leaves, branches and feminine reproductive

Figure 2-a - *Rubidgea* sp. (Papaléo outcrop (lower horizon, Rio Bonito Formation) x 0,5; b - *Gangamopteris* sp., (Acampamento Velho outcrop, Itarare Group) x 1; c - *Rubidgea* sp. (Acampamento Velho outcrop, Itarare Group) x 0,5; d, g, h - *Gangamopteris buriadica* (Acampamento Velho outcrop, Itarare Group) x 1; e - *Glossopteris* sp., evidence of herbivory (roof-shale, Faxinal mine, Rio Bonito Formation) x 0,8; f- *Glossopteris* sp. (Quiteria outcrop, Rio Bonito Formation) x 0,4; i,j - *Glossopteris occidentalis* (roof-shale - Faxinal mine - Rio Bonito Formation) x 1.

Figure 3 - a - *Sphenopteris cf. ischanovensis* (roof-shale, Faxinal mine, Rio Bonito Formation) x 2; b - *Phyllothea indica* (Papaleo outcrop, roof-shale, Rio Bonito Formation) x 0,8; c,f - *Glossopteris communis* (Papalpéo outcrop, roof-shale, lower horizon) x 1- x2; d - *Ginkgoites eximia* (Faxinal outcrop, Itarare Group) x 1,5; e - *Gondwanastachys* sp. (Quiteria outcrop, Rio Bonito Formation) x 2; g, h - *Brasilodendron pedroanum* (Quiteria outcrop, roof-shale) x 0,2; i - *Cordaites* sp. (Papaleo outcrop, roof-shale, upper horizon) x 0,5; j - *Botrychiopsis plantiana* (Papaleo outcrop, lower horizon, Rio Bonito Formation) x 1.

structures of glossopterids constitute 70% of the whole association (*Glossopteris brasiliensis* Guerra-Sommer 1992, *G. papillosa* Guerra-Sommer 1992, *G. similis-intermittens* Guerra-Sommer 1992, *Plumsteadia sennes* Rigby 1963, *Platycardia* sp.), cordaitan leaves (*Ruffloria gondwanensis* Guerra-Sommer 1989) occur as a subordinate group (20%); very delicate filicoid fronds (*Sphenopteris* cf. *ischanovensis*) frequently cross-cutting the lamination represent herbaceous forms that lived under the canopy. Epidermis analyses on cuticles of glossopterid revealed xeromorphic patterns, similar to those from Raniganj coal fields. Studies on recent floras suggest that such features may be related to nitrogen restriction, that occurs nowadays in temperate climate swamps (Lotschert 1969). On the other hand, phytofag evidence in leaves, caused by herbivore insects suggest that the climate was mild. (Guerra-Sommer 1995)

The cordaitan leaves found in this association are assigned on the basis of their epidermic patterns to *Ruffloria* MEYEN, a typical Angara flora form, which is the most important element in the peat-forming plant association in the Late Paleozoic coals in Angara Province (Meyen 1987).

Herbaceous articulates of hygrophytic habitat such as *Phyllothea indica*, and its fertile structures like *Gondwanostachys*, predominate in a megaflora recovered from an antracophylic association at Pantano Grande region near the Leão Mine (Guerra-Sommer *et al.* 1995). Glossopterids of meso-hygrophytic habitat are important forms (*Glossopteris communis*, *G. browniana*, *G. occidentalis* White 1908, *Gangamopteris mosesii* Dolaniti 1954, and *Rubidgea* sp.). Feminine reproductive structures (*Arberia minasica* (White) Rigby 1972) are common elements. Cordaitan leaves (*Cordaite hislopi*) and seeds (*Cordaicarpus irapuensis* Oliveira 1977 and *Samaropsis* sp.) are less frequent. Delicate filicoid fronds (*Sphenopteris* sp., *Rhodopteridium* sp.) occur as complementary elements. Damaged secondary shoots of *Buriadia* sp. are rare in this association.

At the upper section of the Papaleo outcrop (Burjack *et al.* 1982) antracophylic floras associated with two minor coal seams are dominated by arborescent glossopterids with pinnate venation (*Glossopteris communis*, *Glossopteris occidentalis*). Glossopterids with palmate venation (*Rubidgea* sp., *Gangamopteris obovata*) as well as Cordaitales (*Cordaite hislopi*) and Ginkgoales (*Dicranophyllum* and *Cheirophyllum*) are complementary elements. *Phyllothea indica* is a common herbaceous element (Guerra-Sommer *et al.* 1993). Dense associations of horizontally compressed stumps of Lycophyta (*Brasilodendron pedroanum*, Chaloner, Leistikow, Hill 1979) are important elements in the antracophylic association of the upper coal seam. Fronds of *Pecopteris* sp. with fertile structures corresponding to *Asterotheca* sp. were registered in this level by Iannuzzi and Vieira (1997).

On the other hand, the occurrence of impressions of *Cordaite* leaves in barren strata from the yellow sandstones to siltstones suggest that different taxa in the group would be adapted to distinct habitats, in wet and dry biomass.

Arborescent basal stumps of Lycophyta *in situ* cross-cutting the stratification are the dominant higrophyllous element in an

autochthonous antracophylic megafloral association at Quiteria region. The basal portions of these stems related to *Brasilodendron* are depicted as a rounded cornlike structure with roots. Filicoid fronds are also represented, as well as to *Botrychopsis* sp and conifers fragments, representing sterile and fertiles shoots. Casts of lycophytes compressed longitudinally occur in the sequence, parallel to bedding, associated to rare fragments of glossopterids and cordaitan leaves. Jasper & Guerra-Sommer (1998) using taphonomic criteria corroborate the hypothesis of lagoon-barrier system for deposition of the sequence.

Transgressive surface erosion is observed at the top of coalbearing strata. Above this lag there is a package of storm generated sandstones, covered by offshore shales probably corresponding to the maximum flooding surface of the sequence in Southern Brazilian Gondwana (Milani *et al.* in press). In lithostratigraphic terms, the transgressive surface corresponds to the base of Palermo Formation.

The most significant taxa of Early Permian sequences in South Brazilian Gondwana are illustrated in figs. 2 and 3. All material mentioned in this paper is kept at the Palaeobotany section, Institut of Geosciences, UFRGS.

FINAL CONSIDERATIONS Palaeobotanical data allowed to establish a climatic evolution during the deposition of Early Permian sequences in South Brazilian Gondwana. It is, therefore, likely that a changeover occur from a cool cyclic climate at the basal strata to the moist, cyclic interval during coal deposition. This climatic change should be significant to the wetland vegetation which developed in lowlands at the time of the deposition of the peats. Thus, the colonization by arborescent Lycophyta, Filicophyta and herbaceous Sphenophyta indicates not only the ecological conditions but also an adaptation to climatic improvement.

It is important to observe that the paleobotanical data here presented allowed to indicate a mild climate at the time of deposition of roof-shale floras in the Southern Brazilian Gondwana. This assumption is in discordance with the inference of a cold temperate climate suggested before (Rigby 1970, Bernardes de Oliveira 1977), but in accordance to Gastaldo *et al.* (1996) which affirm that the temperate areas of Gondwana throughout the Permian was warmer than hypothesized by climatic models. Comparing Early Permian palaeofloras of Southern Brazilian Gondwana with global ice extent based upon tillites and icrafted deposits becomes clear that the correlation of Sakmarian-Kungurian floras was markedly influenced by the waning of a Permian ice age.

The importance of arborescent lycophyte trunks and stems in roof-shale floras suggested that this group of plants could be important elements in the original biomass of some peat-forming coals in South Brazilian Gondwana. This inference is in accordance with the dominance of lycophyte spores in coal bed assemblages (Marques-Toigo & Correa da Silva 1984) and according with Archangelsky & Cesari (1990) with the linkage between these spores and an arborescent habitat based on ultrastructural studies of exine.

References

- Arber A.N. 1905. *Catalogue of the fossil plants of Glossopteris Flora*. London, Department of Geology British Museum (Natural History), 255 p.
- Bernardes de Oliveira M.E.C. 1977. *Tafoflora Eogondwanica da camada Irupud, Formação Rio Bonito, Grupo Tubarão*. Universidade de São Paulo, São Paulo, Tese de Doutorado, 301 p.
- Burjack M.I. A. 1978. *Estudo Palinológico da Jazida Carbonífera de Charqueadas, Rio Grande do Sul*. - Goiânia, Editora da Universidade de Goiás; 204 p.
- Burjack M.I.A., Klepzig M.C., Fabrício M.E.D., Sommer M.G., Marques-Toigo M., Paim P.S.G., Lavina E.L. 1982. Perfil paleoecológico do afloramento Morro Papaleo, Permiano inferior da bacia do Paraná, RS, Brasil. In: SBG, Congresso Brasileiro de Geologia, 31, Salvador, *Anais*, 4:1260-1270
- Carruthers W. 1869. Coal plants from Brazil. On the Plant Remains from the Brazilian Coal Beds with Remarks on the Genus *Flemingites*. In: Plant N. The Brazilian Coal Fields. *Geological Magazine*, London, 6(4):5-10
- Cazzulo-Klepzig M. & Guerra-Sommer M. 1983. Relationship between the Taphoflora of the Itararé Group, Paraná basin, Southern Brazil and the Permo Carboniferous boundary. In: Dixieme Congres International de Stratigraphie et Geologic du Carbonifere, Madrid, Espanha. *Compte Rendu*, 4:395-402
- Feistmantel O. (1876-1886). The fossil flora of the Gondwana System. Memoirs of the Geological Survey of India. *Paleontologia Indica*, Calcutta, Serie 12,3(1): 1-49
- Gastaldo K.A., DiMichele N.A., Pfefferkorn H.W. (1996). Out of the Icehouse into Greenhouse: a late Paleozoic analog for modern global vegetational change. - *GSA Today*, Boulder, Colorado, 6(10): 1-7.
- Guerra-Sommer M. 1989. Megafloras ocorrentes em horizontes associados a camadas de carvões no Rio Grande do Sul. *Acta Geologica Leopoldensia*, 12 (29):93-104
- Guerra-Sommer M. 1992. Padrões Epidermicos de Glossopteridales da Tafoflora do Faxinal (Formação Rio Bonito Artinskiano/Kunguriano, Bacia do Paraná, Brasil) *Pesquisas*, 1(19):26-40
- Guerra-Sommer M., Marques-Toigo M., Correa da Silva Z.C. 1991. Original biomass and coal deposition in Southern Brazil (Lower Permian, Paraná basin). *Bulletin Soc. Geol. France*. 162(2):227-37
- Guerra-Sommer M. & Cazzulo-Klepzig M. 1993. Biostratigraphy of the Southern Brazilian Neopaleozoic Gondwana Sequence: a preliminary approach. In: XII International Congress of Carboniferous and Permian Stratigraphy and Geology, Buenos Aires, Argentina. *Compte Rendu II*: 61-72.
- Guerra-Sommer M., Marques-Toigo M., Cazzulo-Klepzig M. 1995. Paleoclimatic implication of Lycophyta in the Gondwana of Southern Brazil. *Pesquisas*, 22:21- 31
- Holz M. 1998. The Eopermian Coals of the Paraná basin in Southernmost Brazil: an analysis of the depositional conditions using sequence stratigraphy concepts. *International Journal of Coal Geology*, 36:141-163
- Iannuzzi R. & Vieira C. L. 1997. A presença de Filicopsida no afloramento do Morro Papaleo, município de Mariana Pimentel, RS, Formação Rio Bonito, Eopermiano de Bacia do Paraná. IX RPP, *Revista da Universidade de Guarulhos, Geociências*, n° especial: 224-5
- Jasper A. & Guerra-Sommer M. 1998. Licofitas cormofíticas arborescentes do Afloramento Quiteria, Formação Rio Bonito, Bacia do Paraná. *Pesquisas*. 25:43-60

- Lundqvist G. 1919. Fossile Pflanzen der *Glossopteris* Flora aus Brasilien. *Kungliga Svenska Vetenskaps Akademiens Handlingar*, Stockholm, 60(3): 1-36
- Marques-Toigo M. & Corrêa da Silva Z.C. 1984. On the origin of Gondwanic south brazilian coal measures. *Comun. Serv. Geol. of Portugal*, 70:151-60.
- Mendes J. C. 1952. The Gondwana Formations of Southern Brazil - some of their stratigraphical problems, with emphasis on the fossil Flora. *The Paleobotanist*, 5. Lucknow, 1:335-45
- Meyen S. V. 1987. *Fundamentals of Palaeobotany*. Eds. Chapman & Hall, London, 432 p.
- Milani E. J., Faccini U. F., Scherer C. M. S., Araujo L. M., Cupertino J. A. 1997. Sequences and stratigraphic hierarchy of the Paraná basin (Ordovician to Cretaceous), Southern Brazil. In: Amos A. J., Lopez-Gamundi O., Rocha-Campos A. C. (eds.) - *Sedimentary Basins of South America*, Amsterdam, Elsevier. (in press)
- Pasqualini M., Cunha A. S., Guerra-Sommer M., Piccoli A. E. M. 1986. Análise paleoecológica de seqüências paleoflorísticas na área de Mariana Pimentel - Guaíba, RS. In: SBG, Congresso Brasileiro de Geologia, 34, Goiania. *Anais*, 1:556-569
- Plant V. 1869. The Brazilian Coalfields. *Geological Magazine London*, 6(4):1-10
- Read C. B. 1941. Plantas fósseis do Neopaleozoico do Paraná e Santa Catarina. Monografia. Div. Geol. E Mineralogia, DNPM (12): 1-102
- Rigby J. F. 1970. The distribution of Lower Gondwana plants with the Paraná basin of Brazil. In: Gondwana Symposium, 2, Pretoria, *Proceedings and Papers*, p. 575-84
- Scherer C. M. S. 1999. Análise estratigráfica e litofaciológica da Formação Botucatu (arcabouço da Bacia do Paraná no Rio Grande do Sul) - unpublished PHD Thesis, UFRGS. Porto Alegre, 220p.
- Seward A. C. 1903. Floras of the Past: their composition and distribution. *Nature*, 8:556-68
- White I. D. 1908. Fossil Flora of the Coal Measures of Brazil. In: *Relatório Final da Comissão de Estudos das Minas de Carvão de Pedra do Brasil*. Rio de Janeiro, Imprensa Nacional. p. 1-300.
- Zeiller R. 1895. Note sur Flore Fossile des Gisements Houillers de Rio Grande do Sul (Bresil Meridionale). *Bulletin de la Societe Geologique de France*, 3(23):601-29

Contribution IGC-005

Received January 10, 2000

Accepted for publication May 23, 2000