STUDIES ON THE GEOMORPHIC PROCESSES OF OVERLAND FLOW AND MASS MOVEMENTS IN THE BRAZILIAN GEOMORPHOLOGY

OLGA CRUZ*

Abstract There have been sharp changes in Brazilian geomorphological studies, from the fifties to the seventies of the twentieth century, due to new theoretical developments that have been opposed to the Davisian paradigm. Such studies of deposits correlative to erosional episodes in the Cenozoic stimulated investigation and interpretation with new techniques of weathered, pedogenetic and colluvial materials. They led to constitution of researcher teams under the leadership, among others, of Ab'Saber, Bigarella and Tricart. At the time, other research was already under way on the estimation of losses by erosion in Brazilian agricultural soils.

Geomorphologic research on a broader scale was much influenced, during the seventies, by several external factors, such as techniques of remote sensing, new publications stressing the role of tropical and subtropical process studies, relating forms, materials and processes, catastrophic landslides in São Paulo and Rio de Janeiro, in the 1966/67 summer, new knowledge of quantification, geochemistry, soil mechanics, meteorology and climatology of atmospheric circulation on the South Atlantic, forest clearing, draughts and fight against soil erosion.

The visits of professor Yan De Ploey from the Catholic University of Leuven, Belgium, in 1977/78/79, fostered much Brazilian research on the dynamics of natural geomorphic processes happening on hillslopes, plains and coasts of tropical humid areas, such as overland and channel flow, and mass movements. De Ploey influenced and brought about remarkable changes in several geomorphological trends, always linked to field experimentation, such as the application of troughs to collect pluvial waters on slopes, or studies on the movement of the thick weathered materials and colluvia using sheargraph, or still detection of gullied risk areas and measurements of fluvial waters velocity and transportation power.

In the eighties and nineties, attention to erosional phenomena indicated that natural processes are becoming faster, influenced by anthropic dynamics which interferes with nature. The consolidation of the Brazilian Geomorphology Union confirms the wish for more participation of geomorphologists in the study of contemporary serious environmental problems.

The development of new paradigms in geomorphology and the application of new techniques in the decades between the fifties and the seventies of the XX century, necessitated deep changes in Brazilian geomorphological studies. One slowly passes from a geomorphology of observational research, in regional to continental scale, on Cenozoic erosional levels and surfaces, into more detailed scales, focusing on the study of hillslopes and their correlative deposits.

Methodological orientation from foreign geographers and, specifically, the visits of scientists during the International Geographical Congress in Rio de Janeiro in 1956 decidedly influenced Brazilian Geomorphology. Texts were published, mostly French, on the evolution of the relief, as well as on the effects of climate and vegetation in the diverse morphoclimatic and structural regions of the world. According to Ab'Saber (1958), there was a "true internal revolution in modern Brazilian geomorphology".

Starting in the sixties, there was a strengthening of the studies of correlative deposits related to topographic and planation levels of the Cenozoic erosional phases. Sedimentological research stimulated investigation of depositional features in soil profiles, affording firm bases for the interpretation of weathering, pedogenetic and colluvial materials in erosional levels and on hillslopes. It was a phase of intensive team research under the leadership of, among others, A.N. Ab'Saber, J.J. Bigarella, R. Mousinho and J.X. Silva, G.O. Andrade, G.V. Barbosa, T.C. Silva, A. Christofolletti and A.A.N. Moreira, working on mapping and, specially, on the Quaternary of the country.

At this time, Ab'Saber founded the bulletin *Notícia Geomorfologica*, in São Paulo, later taken over by A. Christofolletti in Campinas. With his extraordinary gift for synthesis and full knowledge of the diverse Brazilian landscapes, Ab'Saber organized geomorphologic and biogeographic maps, present and Quaternary, of the great morphoclimatic domains and phytogeographic provinces of Brazil (Ab'Saber 1966, 1970, 1973). He published his maps in a deeply original and ingenious presentation of Brazilian tropical and subtropical landscapes. Meanwhile he proceeded to research the Quaternary, and established three main levels of geomorphological research: i) compartmentation of relief units according to form and topography; ii) superficial structure and respective materials; iii) physiology and dynamics of present processes (Ab'Saber 1969). Within this methodological orientation, Ab'Saber called attention to the action of geomorphic processes on hillslopes. Ab'Saber already had national stature as a geographer-geomorphologist, as a teacher, researcher and human being; he took his place as a confirmed leader of Brazilian Geomorphology.

João Jose Bigarella pointed towards new research lines on correlative deposits of hilltops and slopes, in diverse Brazilian levels and surfaces molded during the Cenozoic. His texts, illustrated with detailed and enlightening sequential-stratigraphic profiles, afforded regional, local and punctual instances. In 1965, Mousinho and

Bigarella, writing on processes of mass movements in the transportation of debris from rock weathering referred to rainwash action, creep, solifluxion, sliding of coluvia and on their importance to the evolution of hillslopes. Bigarella *et al.* (1970) worked again on sliding, this time on marine sands at Guaratuba, Parana. With his group, Bigarella organized the International Symposium on the Quaternary held in South Brazil in 1975.

Tricart (1961) and Tricart & Cailleux (1965), in papers and conferences on relief forms in tropical humid countries, refer to weathering and pedogenesis, to rainwash and river flow and to mass movements. They assert that a hillslope should not be defined only through its surface and declivity, but also through dynamical aspects of erosion, transport and accumulation in space and time. Such new approaches were particularly well presented by Tricart (1977).

Experiments on soil erosion had been in progress in the U.S.A. since the twenties; also Wischmeier (1959) had made use of the Universal Soil Loss Equation by standard plots. In Brazil, Bertoni (1978), Lombardi and Bertoni (1975) and Lombardi *et al.* (1979) were pioneers, working with field plot experiments at the Instituto Agronomico de Campinas. As early as 1940, De Martonne called attention to mass movements of alteration materials on southeastern Brazilian hillslopes. In 1949, Stenberg referred, at the National Geography Council in Rio de Janeiro, to floods and mass movements in the Paraíba valley. Ruellan (1953) stressed the action of overland flow during rainstorms (*enxurradas*) on the Brazilian relief, calling attention to the strength and impact of tropical rains. Queiroz Neto & Christofolletti (1963) also referred to the importance of rainwash on slopes in the interior of São Paulo, and Rochefort (1958) related rainfall to river flow in tropical and subtropical Brazil.

Other Brazilian and foreign authors continued to stress the role of tropical and subtropical areas to studies of processes, with the help of photo-interpretation techniques. In the 1966-1967 summer, continuous rains, in long and intense episodes on the scarps of the Serra do Mar, in the states of Rio de Janeiro and São Paulo, triggered catastrophic planar landslides and avalanches, with floods, destruction of hydroelectric plants and highways. The need became evident for the study of processes (Meis & Silva 1968, Cruz 1974, 1975, Guidicini *et al.* 1976, Coelho Neto 1979, Bigarella & Suguio 1979).

New analytical studies on hillslopes stressed the relationships between forms, materials and processes. New ideas along the lines of the geosystems paradigm, of quantification data (Christofolletti 1971) and photointerpretation, both of radar (Projeto RADAM) and satellite images (INPE), went together with new meteorological and climatological informations on atmospheric dynamics and South Atlantic cold fronts, later complemented by data on El Niño and La Niña and their pluviometric and hydrologic consequences. New laboratories of geomorphology, pedology, remote sensing and sedimentology were opened consequent to the acquisition of

* Rua Pedroso de Moraes, 684, ap. 602, CEP 05420 - 001, São Paulo, SP

geochemical and soil mechanics information. Additionally, destruction of biodiversity, deforestation, dry periods and fight against erosion have led, in the seventies and eighties, to enhanced interest in research on a broader scale and on the study of the mechanisms of hillslope processes.

Brazil extends from the equatorial to the subtropical belt. The largest part of its eastern façade is characterized by the *mar de morros* (sea of hills) domain (Ab' Saber 1966), covered with Atlantic Forest, with convex upslopes and rectilinear lower slopes, sometimes with benches or shoulders, sometimes concave by the colluvial accumulation, forming *rampas* or debris cones. In the Southeast and partly in the South, the hills are preceded by the scarps of the Serra do Mar, which limits the Atlantic plateau. Three main groups of natural processes take place on the slopes of these humid tropical areas: (i) weathering-pedogenesis, forming thick weathered mantles reworked by (ii) overland and sub-surface flow, and river flow, and by (iii) mass movements, especially on scarped slopes, such as those of Serra do Mar and Serra da Mantiqueira, in the Southeast, and the Serra Geral, in the South. Such natural processes vary according to geographic position, declivity, range and length of the scarps, as well as their geological nature, the thickness of alteration beds and climatic conditions. An increasing and relentless anthropogenic influence has speeded up such dynamics, especially in intensively inhabited areas of risk, at times resulting in catastrophic debris and mudflows.

I.G.U. Commissions in 1976 and 1980, on geomorphological processes and experimental geomorphology, measurements, theory and application in geomorphology, international publications, especially those aiming at processes, as well as Brazilian publications on gullying, runoff and human action have stimulated Brazilian studies on processes. The visits of Professor Yan De Ploey from the Catholic University of Leuven-Belgium encouraged and oriented research on process dynamics, especially fieldwork. De Ploey had been in Brazil in 1975 for the first time, to take part in the International Symposium on the Quaternary. In 1977 he came to the Department of Geography, University of São Paulo, to give an intensive course, to give talks and to participate in fieldwork with postgraduate students. His presence then and again in 1978 and 1979 had an enormous importance by influencing, helping and even changing methods in the sense of stressing field experimentation. Such are the analyses of mass movements in the Serra do Mar scarps (De Ploey & Cruz 1979, De Ploey 1988) and Campos do Jordão Plateau (De Ploey *et al.* 1983), as well as gullying at the limits of the São Paulo sedimentary basin (Cruz & Modenesi 1978). New field techniques were then tried, including the application of sheargraph and rain simulator and measurements of depositional materials from landslides (Cruz *et al.* 1980).

Especially important were the results of a research on the relationship between quantity and intensity of rainfall and the sheetwash with the materials transported, collected, for three years, in Gerlach troughs on scarps under Atlantic Forest, or planted with Pinus, on slide scars (Cruz 1982). Some of the conclusions refer to the importance of herbaceous vegetation and litter layer on the reduction of erosion by sheetwash under forest. Not less important is the fact that troughs on slide scar slopes in reforested area collected a greater load of solid materials than troughs under forest, i.e., there was more erosion on the reforested area.

Postgraduate students followed De Ploey's line producing master dissertations and doctorate theses, such as Baccaro (1983 and 1990), on processes of mass movements on the Serra do Mar scarps and on overland flow in sandstone areas under cerrado in the Brazilian Midwest; Pianaro (1982), on the hydrodynamics of one of the basins of Serra do Mar, in Parana; Domingues (1983) on river flow and hillslope evolution in one of the river basins of the Serra do Mar, in São Paulo; Carvalho (1984) studies the sheetwash with troughs on coastal slopes in São Paulo; Mauro (1990) analyzes gullying in sandstones and basalts on the São Paulo western plateau; Jordão (1991) studies the relationships between mining and hillslope evolution; Furian (1988) studies sheetwash in experimental plots on the Atlantic Plateau and presents a thesis at Caen, France, on the present dynamics of slides, starting from the relationships between morphogenesis and pedogenesis in the evolution of the Serra do Mar slopes; Colangelo (1990) studies mass movements on the lakeside slopes of the Paraiçuna dam, in the Atlantic Plateau. He joins the analysis of balance limit to the model of infinite slope, aiming at an evaluation of mass movements based in data on previous landslides.

At the same time, De Ploey visited the Department of Geosciences at de Federal University of Rio de Janeiro, lecturing and participating in fieldworks organized by Mousinho de Meis. There was an important meeting at Bananal, on gullying and the application of tensiometers and collecting troughs in the study of overland flow in exhausted soils of old coffee plantations on the Precambrian Atlantic Plateau. Under De Ploey's supervision, Coelho Neto (1985) presented a doctor's thesis on soil erosion and hillslope hydrology, in Leuven, Belgium. Guerra (1994) studied soil erosion processes in simulated rainfall experiments and Cunha (1993), fluvial dynamics and drainage basins.

It is important to stress the role of the Brazilian Association of Engineering Geology, which promoted a first symposium at Curitiba in 1980 and a second in 1981 on the control of erosion. Besides discussions of problems of urban and rural erosion, the symposia included contributions on the erosional processes linked to highway systems, concluding that human intervention is the major responsible for the worsening of the erosion processes, due to a total lack of ecological awareness, of the communities, of technicians and of the public administration. One of the data is that agriculture loses, yearly, in the State of São Paulo, 194 million tons of fertile soil. In the 1984 (Belo Horizonte) and 1987 (São Paulo) symposia, special attention was given to erosional processes in Serra do Mar and Serra Geral slopes, to gullying (*voçorocas*), to erosion of reservoir margins, and on their hydrogeologic impact. In 1993, besides these items, other work focused on the increased sediment yield consequent to flooding in urban areas, on the environmental impact of refuse, as well as in cartography, and managing of risk areas. In 1996 and 1999, were discussed diagnoses and proposals for the evaluation and treatment of risk areas, on landfill with solid refuse and on contamination of subsuperficial and subterranean water, on alteration and laterization of soils and on problems stemming from mining activities.

Several works, especially from government agencies, have been presented with agronomical emphasis, trying to relate erosion to rainfall, and specifically to its intensity. Other experimental work has been done on closed plots under diverse plant cover and uses of slope soils. Such experiments perform statistical analysis of superficial materials, or measurements of creep, or of rain erosivity, sometimes applying the Bertoni (1978) equation, based in Wischmeier. Such are the theses by Casseti (1984) on the Goiania plateau, Pacheco (1991) on the scarps of the Serra Geral in Rio Grande do Sul, Suertegaray (1987) in the sand wastes of Quaraí, Rio Grande do Sul, or still Tavares (1986) in the western plateau of São Paulo. Otherwise, Colangelo (1995) related indexes of rain frequency and magnitude with the hydraulic conductivity of weathered and colluvial materials from slumps on the Atlantic Plateau, and their resistance to shearing, as measured by the sheargraph.

In 1982, Ab'Saber identified critical areas of degradation; he called attention in 1985 to breaks of ecological balance in the Serra do Mar at Paranapiacaba, São Paulo, polluted for at least 20 years by factories in Baixada Santista. Government agencies started strategies of recuperation, with transplantation of native trees and seedlings from the air in areas hard to reach (Pompeia *et al.* 1989). However many irregular dwellings were built, invading other areas identified as of risk. Every summer, in general very rainy, serious slide problems afflict a poor population that ends up by inhabiting shanties or small brick houses in forbidden areas. This leads to impermeability and weakening of the slopes, robbing them from their natural defenses and causing landslides. It is important to note that such areas of risk are in general characterized by elongate steep slopes and broad topographical ranges, or, on the plains, connected to meander belts. During very wet summers, with long and intense rainy episodes soils and thick bedrock weathered materials, with many blocks, tend to move, even under forest or in areas of reforestation. They are of course areas of risk and of permanent protection.

The accentuation of current erosional processes to be seen on several sectors of the Brazilian coast, especially in the last two decades, has promoted studies on wave and tidal currents and on the influence of climatic processes on the beach and foreshore. Tide measurements by oceanographic institutions, application of geoprocessing techniques and experimental research on littoral and coastal areas were made in São Paulo by Modenesi (1969), Furtado (1978), Batres (1978), Modenesi *et al.* (1983), Poncano (1985), Tessler and Furtado (1983), Souza (1997); in Rio Grande do Sul, by Villwock (1987), Bernardi *et al.* (1987), Tomazelli & Vilwock (1989); in Rio de Janeiro, by Muehe

(1984), Muehe *et al.* (1989); in Bahia, by Farias *et al.* (1985), Bittencourt *et al.* (1990); in Maranhão, by Feitosa (1996); in Santa Catarina, by Duarte (1981, 1995), Abreu Castilhos & Correa (1991), Horn Filho *et al.* (1994), Ore *et al.* (1994), Pellerin & Paulino (1996), Diehl (1997), Cruz (1998), Herrmann (1999); in Paraná, by Bigarella (1978), Angulo (1989), Kohler (1996), Canali *et al.* (1999), among many Brazilian researchers, in seashore coastal areas and in neighboring countries, as for example, in the French Guiana (Prost 1986).

The Committee on Geomorphology and Environmental Change of I.G.U., new publications on tropical geomorphology (Thomas 1994)

References

- Abreu Castilhos J.A. & Correa C.H.T. 1991. Avaliação preliminar dos processos erosivos na praia da Armação - Ilha de Santa Catarina, Florianópolis-SC. In: Simp. Geogr. Física Aplicada, 4, *Anais*, 1:218-229
- Ab'Saber A.N. 1958. A Geomorfologia no Brasil. São Paulo-SP, USP, *Notícia Geomorfologica*, 2:1-8
- Ab'Saber A.N. 1966. *O domínio das "mares de morms" no Brasil*. São Paulo-SP, IGEOG-USP, 9 p. (Geomorfologia 2)
- Ab'Saber A.N. 1969. *Um conceito de geomorfologia a serviço das pesquisas sobre o Quaternário*. São Paulo-SP, IGEOG-USP, 23p. (Geomorfologia 18)
- Ab'Saber A.N. 1970. *Províncias geológicas e domínios morfoclimáticos no Brasil*. São Paulo-SP, IGEOG-USP, 26p. (Geomorfologia 20)
- Ab'Saber A.N. 1973. *A organização natural das paisagens inter e subtropicais brasileiras*. São Paulo-SP, IGEOG-USP, 14 p. (Geomorfologia 41)
- Ab'Saber A.N. 1982. *Degradação da natureza por processos antrópicos, na visão dos geógrafos*. São José do Rio Preto-SP, IBILCE-UNESP, 27p. (Inter-Facies 106)
- Ab'Saber A.N. 1985. A gestão do espaço natural (relembrando Caraguatubá 1967, para compreender Cubatão, 1985). *Rev. Arquitetura e Urbanismo*, 1:90-93
- Angulo R.J. 1989. Variações na configuração da linha de costa do Paraná nas últimas quatro décadas. In: ABEQUA, Congr., 2, Rio de Janeiro-RJ, *Public Especial*, 1, p.24
- Baccaro C.A.D. 1983. *Os processos de movimentos de massa e a evolução das vertentes na serra do Mar*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 165p.
- Baccaro C.A.D. 1990. *Estudo dos processos geomorfológicos de escoamento pluvial em área de cerrado, Uberlândia-MG*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 164p.
- Batres V.B.K. 1978. *A foz do rio Ianhema e áreas adjacentes, estudo de geomorfologia costeira*. Dep. de Geografia - FFLCH - Universidade de São Paulo-SP, Dissertação de Mestrado.
- Bernardi H., Cordazzo C.V., Costa C.S.B. 1987. Efeito de ressacas sobre Blutaparon portulacoides (St. Hill) Mears nas dunas costeiras do sul do Brasil. São Paulo-SP, *Cienc. e Cultura*, 39:345-347
- Bertoni J. 1978. Nomograma para solução de perdas do solo do Estado de São Paulo. In: II Encontro Nacional de Pesquisa para Conservação do Solo, Passo Fundo-RS, *Anais*
- Bigarella J.J. 1978. *A Serra do Mar e a porção oriental do Estado do Paraná*. Curitiba, Secretaria do Planejamento and ADEA, 249 p.
- Bigarella J.J., Silva J.X., Duarte G.M. 1970. O desastre de Guaratuba-PR. Curitiba-PR, *Rev. IBPT*, 14
- Bigarella J.J. & Suguio K. 1979. *Ambiente fluvial: ambientes de sedimentação, sua interpretação e importância*. Curitiba, ADEA, 183p.
- Bittencourt A.C.S.P., Farias F.F., Boas G.S.V. 1990. Influência da deriva litorânea no desenvolvimento do esporão de Caixa-Pregos, baía de Todos os Santos-BA. São Paulo-SP, *Rev. Bras. de Geociências*, 20:197-200
- Canali N.E., Oka-Fiori C., Guedes J.A., Nunes F.G. 1999. The Antonina and Paranaguá way's region, Brazil: environmental zoning. In: Regional Conference Geomorphology, Rio de Janeiro-RJ, *Abstracts*, p.24
- Carvalho G.R.F. 1984. *Contribuição ao estudo da dinâmica atual geomorfológica do morro da Península em Guarujá-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 147p.
- Gassed V. 1984. *Estudo dos efeitos morfodinâmicos pluviais no planalto de Goianina-GO*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 138p.
- Christofolletti A. 1971. *A teoria dos sitemas*. Bol. de Geografia Teórica, 1:43-60
- Coelho Neto A.L. 1979. *O processo erosivo nas encostas do maciço da Tijuca-RJ*. Inst. de Geociências, Universidade Federal do Rio de Janeiro, Rio de Janeiro-RJ, Dissertação de Mestrado
- Coelho Neto A.L. 1985. *Surface hidrology and soil erosion in tropical mountainous rainforest drainage basin Rio de Janeiro*. Inst. de Geociências, Universidade Federal do Rio de Janeiro, Rio de Janeiro-RJ, Tese de Doutorado
- Colangelo A.C. 1990. *Movimentos de massa e evolução geomorfológica nas vertentes marginais do lago-barragem de Paraibuna-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado
- Colangelo A.C. 1995. *Teste de experimentação com "sheargraph" e obtenção de limiares nas vertentes marginais do lago-barragem de Paraibuna-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado
- Cruz O. 1974. *A serra do Mar e o litoral na área de Caraguatubá-SP*. Inst. de Geografia, Universidade de São Paulo, São Paulo-SP, 181 p. (Serie Teses e Monografias, 11)
- Cruz O. 1975. Evolução de vertentes nas escarpas da serra do Mar em Caraguatubá-SP. In: Simpósio Internacional sobre o Quaternário, *Anais*, Academia Brasileira de Ciências-suplemento 47:479-480
- Cruz O. 1982. *Estudos dos processos geomorfológicos do escoamento pluvial na área de Caraguatubá-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Livre Docência, 151p.
- Cruz O. 1998. *A Ilha de Santa Catarina e o Continente próximo. Um estudo de geomorfologia Costeira*. Florianópolis-SC, Editora Universidade Federal de Santa Catarina, 279p.
- Cruz O. & Modenesi M.C. 1978. Some observations on the evolution of a gully in sediments of the São Paulo Basin, São Paulo. In: International Congress INQUA, Birmingham-UK, *Abstracts*, 99:77-85
- Cruz O., Suguio K., Eichler B.B. 1980. Sedimentação acelerada de pontais e barras arenosas na enseada de Caraguatubá-SP. Causas naturais e artificiais. São Paulo-SP, USP, *Bol. Inst. Oceanográfico*, 33(1):39-53
- Cunha S.B. 1993. Impactos das obras de engenharia sobre o ambiente biofísico da bacia do rio São João (Rio de Janeiro-Brasil). Departamento de Geografia. Universidade clássica de Lisboa, Lisboa, 415p.
- De Martonne E. 1940. Problemes morphologiques du Bresil tropical atlantique. *Annales de Geographie*, 49:1-27 et 105-129
- De Ploey Y. 1988. Concerning factors of soil erosion, with special reference to tropical areas. In: Latin American Regional Conference, Rio de Janeiro-RJ, *Abstracts*: 317-322
- De Ploey Y. & Cruz O. 1979. Landslides in the serra do Mar, Brazil. *Catena*, 6:111 -122
- De Ploey Y., Cruz O., Modenesi M.C. 1983. Resistance au cisaillement et conditions de glissements de terrain a Caraguatubá e a Campos do Jordão-SP. In: Estudo e cartografia de formações superficiais e suas aplicações em regiões tropicais. *Colóquio interdisciplinar franco-brasileiro*, Dep. de Geografia, FFLCH, Universidade de São Paulo, São Paulo, 1:393-402
- Diehl F.L. 1997. *Aspectos geoevolutivos, morfodinâmicos e ambientais do pontal da Daniela, Ilha de Santa Catarina*. Dep. de Geociências - CFH - Universidade Federal de Santa Catarina, Florianópolis-SC, Dissertação de Mestrado
- Domingues E.N. 1983. *Estudo de processos geomorfológicos do escoamento fluvial e evolução de vertentes na serra de Cubatão-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 153p.
- Duarte G.M. 1981. *Estratigrafia e evolução do Quaternário no plano costeiro none da Una de Santa Catarina*. Departamento de Geografia, Universidade Federal do Rio Grande do Sul, Porto Alegre, Dissertação de Mestrado
- Duarte G.M. 1995. *Depositos cenozoicos costeiros e a morfologia do extremo sul de Santa Catarina*. Instituto de Geociências, Universidade de São Paulo, São Paulo, Tese de Doutorado
- Farias E.F., Bittencourt A.C.S.P., Zanini Jr. A., Dominguez J.M.L. 1985. Variações temporais e espaciais na dinâmica de sedimentação da praia de Armação, Salvador-BA. *Rev. Bras. de Geociências*, 15:48-54
- Feitosa A.C. 1996. *Dinâmica dos processos geomorfológicos da área costeira a nordeste da ilha do Maranhão-MA*. Inst. de Geociências - Universidade Estadual Paulista, Rio Claro-SP, Tese de Doutorado, 249p.
- Furian S.M. 1994. *Morphogenese/pedogenese em milieu tropical humide de la serra do Mar, Bresil, contribution de l'alteration et de la pedogenese a une dynamique actuelle de glissement*. Caen-France, SLSC, Tese de Doutorado, 178p.
- Furian Dias S.M. 1988. *Os processos geomorfológicos do escoamento pluvial na estacão experimental de Cunha-SP - Um esboço metodológico*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 187p.
- Furtado V. V. 1978. *Contribuição ao estudo da sedimentação atual no Canal de São Sebastião-SP*. Inst. Oceanográfico, Universidade de São Paulo, São Paulo-SP, Tese de Doutorado
- Gr6 J.C.R., Abreu Castilhos J., Horn P.N.O. 1994. Morphodynamic and sedimentological study of the "Mole beach", Santa Catarina island, Brazil. In: Internacional Sedimentological Congress, 14, Recife-PE, *Abstracts*
- Guerra A.J.T. 1994. The effect of organic matter content on soil erosion in simulated rainfall experiments in W. Sussex, U.K. Soil Use and Management, Harpenden.
- Guidicini G. & Iwasa O.Y. 1976. *Ensaio de correlação entre pluviosidade e correções no meio tropical úmido*. Inst. de Pesquisas Tecnológicas, São Paulo-SP, 48p., (Publicação 1080)
- Herrmann M.L.P. 1999. *Estudo de geomorfologia costeira nos municípios de São Jose, Palhoça e Biguaçu-SC*. Universidade Federal de Santa Catarina, Florianópolis-SC, Tese de Doutorado
- Horn Filho N.O., Abreu Castilhos J., Gr6 J.C.R., Diehl F.L. 1994. Estudo geológico correlativo entre o Quaternário costeiro das ilhas de São Francisco do Sul e de Santa Catarina-SC. In: SBG, Congresso Brasileiro de Geologia, 38, Balneário Camboriú-SC
- Jordão S. 1991. *A mineração nos municípios de Guarulhos e Mairiporã-SP. Aspectos geomorfológicos aplicados a sua organização e desenvolvimento*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 250p.
- Kohler V.B. 1996. *De Pontal do Sul a praia de Leste-PR, um estudo de geomorfologia costeira*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado
- Lombardi F.N. & Bertoni J. 1975. *Tolerancia de perda da terra para solos do Estado de São Paulo*. Campinas-SP, Inst. Agrônomo, 12 p. (Bol. Técnico 28)
- Lombardi F.N. *et al.* 1979. *Simulador de chuva e sua utilização em pesquisa de erosão do solo*. Campinas-SP, p. 81-98, (Bol. Inst. Agrônomo 31)
- Mauro C.A. 1990. *Voçorocas: marcas das relevoes sociedade-natureza na bacia do Monjolinho, São Carlos-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 222p.
- Meis M.R.M. & Silva J.X. 1968. Considerações geomorfológicas a propósito dos movimentos de massa ocorridos no Rio de Janeiro-RJ. *Rev. Bras. de Geografia*, 30(1):55-73
- Modenesi M.C. 1969. *Memoria explicativa da carta geomorfológica da ilha de Santo Amaro - SP. Primeiros estudos*. São Paulo-SP, IGEOG-USP, 15 p. (Aerofotogeografia 2)

- Modenesi M.C., Tessler M.G., Cruz O., Coimbra A.M. 1983. Influence of marine and continental processes on the dynamics of a sand-ridge at the mouth of the Massaguaçu river, Caraguatatuba-SP. Preliminary conditions. São Paulo-SP, USP, *Bol. Inst. Oceanográfico*, 32(1):77-81.
- Mousinho M.R. & Bigarella J.J. 1965. Movimentos de massa no transporte dos detritos da meteorização das rochas. Curitiba-PR, UFPR, *Bol. Paranaense de Geografia*, 16 e 17:43-84.
- Muehe D. 1984. Evidências de recuo dos cordões litorâneos em direção ao continente no litoral do Rio de Janeiro. In: Simpósio sobre Restingas, Origem, Estrutura, Processos, Niterói-RJ, *Anais*.
- Muehe D., Correa H.T., Ignaras S. 1989. Avaliação dos riscos de erosão dos cordões litorâneos entre Niterói e Cabo Frio-RJ. In: *Simpósio de Geografia Física Aplicada, Nova Friburgo-RJ*, 1:368-383.
- Pacheco M.M.S. 1991. *Evolução e dinâmica geomorfológica das vertentes na área da escarpa entre Taquara e São Francisco de Paula-RS*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 126p.
- Pellerin J. & Paulino L.A. 1996. Aspectos geomorfológicos da ilha de Santa Catarina. In: SBPC, Reuniao Especial, 3, Florianópolis, *Anais*.
- Pianaro A. 1982. *Estudo hidrodinâmico da bacia do rio Tagaça, Guaraquecaba-PR*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Dissertação de Mestrado, 126p.
- Pompeia S.L., Pradella D.Z.A., Martins S.E., Santos R.S., Diniz K.M. 1989. A sementeira aérea na serra do Mar, Cubatão-SP. *Rev. CETESB de Tecnologia*, 3:13-19.
- Ponçano W.L. 1985. *Sedimentação atual aplicada a portos no Brasil*. Inst. de Geociências, Universidade de São Paulo, São Paulo-SP, Tese de Doutorado.
- Prost M.T. 1986. Aspects of the morphosedimentary evolution of French Guyane Coastline. In: *Quaternary of South America and Antarctic peninsula*, São Paulo-SP, 4:191-218.
- Queiroz Neto J.P. & Christofoletti A. 1963. Ação do escoamento superficial das águas pluviais na serra de Santana-SP. São Paulo-SP, AGB, *Bol. Paulista de Geografia*, 45:59-71.
- Rochefort M. 1958. *Rapports entre la pluviosité et le coulement dans le Brésil subtropical et le Brésil tropical atlantique*. Paris, France.
- Ruellan F. 1953. O Papel das enxurradas no modelado do relevo brasileiro. São Paulo-SP, AGB, *Bol. de Geografia*, 13 e 14.
- Souza C.R.G. 1997. *As células de deriva litorânea e a erosão nas praias do Estado de São Paulo*. Inst. de Geociências, Universidade de São Paulo, São Paulo-SP, Tese de Doutorado.
- Sternberg H.O.R. 1949. Enchentes e movimentos coletivos no vale do Paraíba em dezembro de 1949. Rio de Janeiro-RJ, IBGE, *Rev. Bras. de Geografia*, 11:223-261.
- Suertegaray D.M.A. 1987. *A trajetória da natureza: um estudo geomorfológico sobre os areais de Quaraí-RS*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 243p.
- Tavares A.C. 1986. *A erosão dos solos no contexto da análise ambiental: o exemplo do alto curso do rio São José dos Dourados-SP*. Dep. de Geografia - FFLCH - Universidade de São Paulo, São Paulo-SP, Tese de Doutorado, 254p.
- Tessler M.G. & Furtado V.V. 1983. Dinâmica de sedimentação das feições de assoreamento da região lagunar Cananã/Itaipu-SP. São Paulo-SP, USP, *Bol. Inst. Oceanográfico*, 32:117-124.
- Thomas M.F. 1994. *Geomorphology in the tropics. A study of weathering and denudation in low latitudes*. J. Wiley & Sons.
- Tomazelli L.J. & Villwock J.A. 1989. Processos erosivos atuais na costa do Rio Grande do Sul, evidências de uma tendência contemporânea de elevação do nível relativo do mar. In: 2º Congresso ABEQUA, Rio de Janeiro-RJ, *Publ. Especial*, 1, 16p.
- Tricart J. 1961. Les caractéristiques fondamentales du système morphogénétique des pays tropicaux humides. Paris, *Information Géographique*, 25:155-169.
- Tricart J. 1977. *Ecodinâmica*, Rio de Janeiro-RJ, IBGE.
- Tricart J. & Cailleux, A. 1965. *Traité de géomorphologie. Le modelé des régions chaudes, forêts et savanes*. Paris, SEDES.
- Villwock J.A. 1987. Processos costeiros e a formação das praias arenosas e campos de dunas na costa Sul e Sudeste brasileira. In: *Simpósio sobre ecossistemas Sul e Sudeste brasileiro, síntese dos conhecimentos*, 1:380-398.
- Wischmeier W.H. 1959. A rainfall erosion index for a universal soil loss equation. In: Soil Science Society of America, *Proceedings*, 23(2):246-249.

Contribution IGC-030
Received January 29, 2000
Accepted for publication April 28, 2000